CURRICULUM VITAE KATIE A. MEYER

Work Address

Nutrition Department University of North Carolina at Chapel Hill 123 West Franklin Street Chapel Hill, NC 27516

Email: ktmeyer@unc.edu

EDUCATION

1997-2002 ScD, Epidemiology, Harvard School of Public Health, Boston, MA.

1995-1997 MPH, Epidemiology, University of Minnesota, School of Public Health, Minneapolis, MN.

1989-1993 BA, Art History, Macalester College, Saint Paul, MN.

WORK EXPERIENCE

2012-present Research Assistant Professor

Nutrition Department

University of North Carolina at Chapel Hill

Chapel Hill, NC

2009-2012 NHLBI CVD Epidemiology Post-Doctoral Research Fellow

Division of Epidemiology and Community Health University of Minnesota School of Public Health

Minneapolis, MN

2002-2009 Chronic Disease Epidemiologist

Division of Chronic Disease Maine Bureau of Health

Augusta, ME

1997-2001 NIH Pre-Doctoral Research Fellow

Harvard School of Public Health

Boston, MA

PUBLICATIONS

Meyer KA, Sijtsma FPC, Nettleton JA, Steffen LM, Van Horn L, Shikany JM, Gross MD, Mursu J, Traber MG, Jacobs DR. Dietary patterns are associated with plasma F₂-isoprostanes in an observational cohort study of adults. Free Radic Biol Med 2012

Sijtsma FP, Meyer KA, Steffen LM, Shikany JM, Van Horn L, Harnack L, Kromhout D, Jacobs DR Jr. Longitudinal trends in diet and effects of sex, race, and education on dietary quality score change: the Coronary Artery Risk Development in Young Adults study. Am J Clin Nutr. 2012 Mar;95(3):580-6. Epub 2012 Feb 1.

Meyer KA, Wall MM, Larson NI, Laska MN, Neumark-Sztainer D. Sleep duration and BMI in a sample of young adults. Obesity (Silver Spring). 2012 Jan 26. [Epub ahead of print]

Jacobs DR, Mursu J, Meyer KA. Editorial: The importance of food. Arch Pediatr Adolesc Med. 2012

Meyer KA, Demerath EW, Hannan PJ, Neumark-Sztainer D. Body fat is differentially related to body mass index in U.S.-born African-American and East African immigrant girls. Am J Hum Biol 2011;23:720-3.

Thyagarajan B, Meyer KA, Smith L, Williams DO, Beckett WS, Gross MD, Jacobs DR. Serum carotenoid concentrations predict evolution of lung function in young adults: the CARDIA study. Am J Clin Nutr Sept 2011 Epub ahead of print.

Meyer KA, Friend S, Hannan PJ, Himes JH, Demerath EW, Neumark-Sztainer D. Ethnic variation in body composition assessment in a sample of adolescent girls. Int J Pediatr Obes 2011;6:481-90.

Jacobs DR, Meyer KA. Invited commentary: On population subgroups, mathematics, and interventions. Am J Epidemiol 2011 Jan 17.

Decker K, Meyer K, Thompson D, Littlefield D. Similar asthma prevalence estimates obtained from pre-adolescent and parent survey responses. J Clin Epidemiol 2008;61:611-6.

Meyer KA, Decker K, Mervis CA, Louder D, Bradshaw J, DeVader S, Wigand D. Emergency medical services data for cardiovascular disease surveillance, program planning, and evaluation in Maine. Prev Chronic Dis 2008;5:A54.

Tippy K, Meyer K, Aronson R, Wall T. Characteristics of coordinated ongoing comprehensive care within a medical home in Maine. Maternal and Child Health J 2005;9(2 Suppl):S13-21.

Meyer KA, Williams P, Hernandez-Diaz S, Cnattingius S. Smoking and the risk of oral clefts: exploring the impact of study designs. Epidemiology 2004;15:671-8.

Meyer KA, Arduino JM, Santanello NC, Knorr B, Bisgaard H. Response to montelukast sodium among subgroups of children aged 2-14 years with asthma. J Allergy Clin Immunol. 2003;111:757-62.

Hernandez-Diaz S, Hernan MA, Meyer KA, Werler MM, Mitchell AA. Case-crossover and case-time-control designs in birth defects epidemiology. Am J Epidemiol. 2003;158:385-91.

Meyer KA, Werler MM, Hayes C, Mitchell AA. Low maternal alcohol consumption during pregnancy and oral clefts in offspring: The Slone Birth Defects Study. Birth Defects Res Part A Clin Mol Teratol. 2003;67:509-14.

Meyer KA, Conigrave KM, Chu NF, Rafai N, Spiegelman DM, Stampfer MJ, Rimm EB. Alcohol consumption patterns and HbA1c, C-peptide, and insulin concentrations in men. J Am Coll Nutr 2003;22:185-94.

Meyer KA, Kushi LH, Jacobs DR Jr, Folsom AR. Dietary fat and incidence of type 2 diabetes mellitus in older Iowa women. Diabetes Care 2001; 24:1528-35.

Jacobs DR Jr, Pereira MA, Meyer KA, Kushi LH. Fiber from whole grains, but not refined grains, is inversely associated with all-cause mortality in older women: the Iowa Women's Health Study. J Am Coll Nutr 2000; 19:326S-330S.

Meyer KA, Kushi LH, Jacobs DR Jr, Slavin J, Sellers TA, Folsom AR. Carbohydrates, dietary fiber, and incident type 2 diabetes mellitus in older women. Am J Clin Nutr 2000; 71:921-930.

Kushi LH, Meyer KA, Jacobs DR Jr. Cereals, legumes, and chronic disease risk reduction: evidence from epidemiologic studies. Am J Clin Nutr 1999; 70:451S-458S.

Yochum L, Kushi LH, Meyer K, Folsom AR. Dietary flavonoid intake and risk of cardiovascular disease in postmenopausal women. Am J Epidemiol 1999;149:943-949.

Jacobs DR Jr, Meyer KA, Kushi LH, Folsom AR. Is whole grain intake associated with reduced total and cause-specific death rates in older women? The Iowa Women's Health Study. Am J Public Health 1999;89:322-329.

Bostick RM, Kushi LH, Wu Y, Meyer KA, Sellers TA, Folsom AR. Relation of calcium, vitamin D, and dairy food intake to ischemic heart disease mortality among postmenopausal women. Am J Epidemiol 1999;149:151-161.

Jacobs DR Jr, Meyer KA, Kushi LH, Folsom AR. Whole-grain intake may reduce the risk of ischemic heart disease death in postmenopausal women: the Iowa Women's Health Study. Am J Clin Nutr 1998;68:248-257.